

Acceleration of the Universe in Type-0 Non-Critical Strings

G. A. Diamandis and **B. C. Georgalas**

*Physics Department, Nuclear and Particle Physics Section, University of Athens,
Panepistimioupolis GR 157 71, Ilisia, Athens, Greece.*

N. E. Mavromatos

*Department of Physics, Theoretical Physics, King's College London,
Strand, London WC2R 2LS, United Kingdom.*

and

E. Papantonopoulos

*Department of Physics, National Technical University of Athens,
Zografou Campus GR 157 80, Athens, Greece.*

Abstract

Presently there is preliminary observational evidence that the cosmological constant might be non zero, and hence that our Universe is eternally accelerating (de Sitter). This poses fundamental problems for string theory, since a scattering matrix is not well defined in such Universes. In a previous paper we have presented a model, based on (non-equilibrium) non-critical strings, which is characterized by eventual “graceful” exit from a de Sitter phase. The model is based on a type-0 string theory, involving D3 brane worlds, whose initial quantum fluctuations induce the non criticality. We argue in this article that this model is compatible with the current observations. A crucial rôle for the correct “phenomenology” of the model is played by the relative magnitude of the flux of the five form of the type 0 string to the size of five of the extra dimensions, transverse to the direction of the flux-field. We do not claim, at this stage at least, that this model is a realistic physical model for the Universe, but we find it interesting that the model cannot be ruled out immediately, at least on phenomenological grounds.

1. Introduction

Recently there is some preliminary experimental evidence from type Ia supernovae data [1], which supports the fact that our Universe *accelerates* at present: distant supernovae (redshifts $z \sim 1$) data indicate a slower rate of expansion, as compared with that inferred from data pertaining to nearby supernovae. Distant supernovae look dimmer than they should be, if the expansion rate of the Universe would be constant.

This could be a consequence of a non-zero cosmological constant, which would imply that our Universe would be eternally accelerating (de Sitter), according to standard cosmology [2]. Such evidence is still far from being confirmed, but it is reinforced by combining these data with Cosmic Microwave background (CMB) data (first acoustic peak) implying a spatially $\Omega_{\text{total}} = 1.0 \pm 0.1$ flat Universe [3].

Let us review briefly the current situation. Specifically, the best fit spatially flat Universe to the data of [1, 3] implies, to 3σ confidence level, that

$$\Omega_{M,0}(\text{matter}) \simeq 0.3, \quad \text{and} \quad \Omega_{X,0}(\text{dark energy}) \simeq 1 - \Omega_M \simeq 0.7, \quad (1.1)$$

where the subscript 0 indicate present values. If the data have been interpreted right this means that 70% of the present energy density of the Universe consists of an unknown substance (“*dark energy*”). For the fit of [1, 3] the dark energy has been taken to be the standard cosmological constant.

An important phenomenological parameter, which is of particular interest to astrophysicists is the *deceleration parameter* q of the Universe [2], which is defined as:

$$q = -\frac{(d^2 a_E / dt_E^2) a_E}{(da_E / dt_E)^2} \quad (1.2)$$

where $a_E(t)$ is the Robertson-Walker scale factor of the Universe, and the subscript E denotes quantities computed in the so-called Einstein frame, that is where the gravity action has the canonical Einstein form as far as the scalar curvature term is concerned. This

distinction is relevant in string-inspired effective theories with four-dimensional Brans-Dicke type scalars, such as dilatons, which will be dealing with here.

It should be mentioned that in standard Robertson-Walker cosmologies with matter the deceleration parameter can be expressed in terms of the matter and vacuum (cosmological constant) energy densities, Ω_M and Ω_Λ respectively, as follows:

$$q = \frac{1}{2}\Omega_M - \Omega_\Lambda \quad (1.3)$$

For the best fit Universe (1.1) one can then infer a present deceleration parameter $q_0 = -0.55 < 0$, indicating that the Universe accelerates today.

If the data have been interpreted right, then there are three possible explanations [2]:

(i) Einstein's General Relativity is incorrect, and hence Friedman's cosmological solution as well. This is unlikely, given the success of General Relativity and of the Standard Cosmological Model in explaining a plethora of other issues.

(ii) the 'observed' dark energy and the acceleration of the Universe are due to an 'honest' cosmological *constant* Λ in Einstein-Friedman-Robertson-Walker cosmological model. This is the case of the best fit Universe (1.1) which matches the supernova and CMB data. In that case one is facing the problem of eternal acceleration, for the following reason: let $\rho_M \propto a^{-3}$ the matter density in the Universe, with $a(t)$ the Robertson-Walker scale factor, and t the cosmological observer (co-moving) frame time. The vacuum energy density, due to Λ , is assumed to be constant in time, $\rho_\Lambda = \text{const}$. Hence in conventional Friedmann cosmologies one has:

$$\Omega_\Lambda/\Omega_M = \rho_\Lambda/\rho_M \propto a(t)^3, \quad (1.4)$$

and hence eventually the vacuum energy density component will dominate over matter. From Friedman's equations then, one observes that the Universe will eventually enter a de-Sitter phase, in which $a(t) \sim e^{\sqrt{\frac{8\pi G_N}{3}}\Lambda t}$, where G_N is the gravitational (Newton's) constant.

This implies eternal expansion and acceleration, and most importantly the presence of a *cosmological horizon*

$$\delta = a(t) \int_{t_0}^{\infty} \frac{cdt}{a(t)} < \infty \quad (1.5)$$

It is this last feature in de Sitter Universes that presents problems in defining proper *asymptotic states*, and thus a consistent scattering matrix for field theory in such backgrounds [4]. The analogy of such global horizons with microscopic or macroscopic black hole horizons in this respect is evident, the important physical difference, however, being that in the cosmological de Sitter case the observer lives “inside” the horizon, in contrast to the black hole case.

Such eternal-acceleration Universes are, therefore, bad news for critical string theory [4], due to the fact that strings are by definition theories of on-shell S -matrix and hence, as such, can only accommodate backgrounds consistent with the existence of the latter.

(iii) the ‘observed effects’ are due to the existence of a *quintessence* field φ , which has not yet relaxed in its absolute minimum (ground state), given that the relaxation time is longer than the age of the Universe. Thus we are still in a non-equilibrium situation, relaxing to equilibrium gradually. In this drastic explanation, the vacuum energy density, due to the potential of the field φ will be time-dependent. In fact the data point towards a $1/t^2$ relaxation, with $t \geq 10^{60}$ in Planck units, where the latter number represents the age of the observed Universe.

It is this third possibility that we have attempted to adopt in a proper non-critical string theory framework in ref. [9]. Non-critical strings can be viewed as non-equilibrium systems in string theory [7]. The advantage of this non-equilibrium situation lies on the possibility of an eventual exit from the de Sitter phase, which would allow proper definition of a field-theory scattering matrix, thus avoiding the problem of eternal horizons mentioned previously. This would be a welcome fact from the point of view of string theory.

However, it should be stressed that our studies, although encouraging, are far from

being complete from a mathematical point of view. Non-critical (Liouville) string theory, with time-like backgrounds, like the ones used in [9], is not completely understood at present [5, 6], although significant progress has been made towards this direction. The main reason for this is the fact that world-sheet Liouville correlation functions, do not admit a clear interpretation as ordinary scattering (S) matrix on-shell amplitudes in target space. Rather, they are S -matrix amplitudes, connecting asymptotic density matrices, as appropriate for the ‘open system’ character of such string theories [7, 8]. Such systems may thus be of relevance to cosmology, especially after the above mentioned recent experimental evidence on a current era acceleration of the Universe.

Exit from de Sitter inflationary phases is another feature that cannot be accommodated (at least to date) within the context of critical strings [7]. This is mainly due to the fact that such a possibility requires time-dependent backgrounds in string theory which are also not well understood. On the other hand there is sufficient evidence that such a ‘graceful exit possibility’ from the inflationary phase can be realized in non-critical strings, with a time-like signature of the Liouville field, which thus plays the rôle of a Robertson-Walker comoving-frame time [7]. The evidence came first from toy two-dimensional specific models [10], and recently was extended [9] to four-dimensional models based on the so-called type-0 non-supersymmetric strings [11]. The latter string theory has four-dimensional brane worlds, whose fluctuations have been argued in [9] to lead to super criticality of the underlying string theory, necessitating Liouville dressing with a Liouville mode of time-like signature. In general, Liouville strings become critical strings after such a dressing procedure, but in one target-space dimension higher. However in our approach [7, 9], instead of increasing the initial number of target space dimensions ($d = 10$ for type-0 strings), we have identified the world-sheet zero mode of the Liouville field with the (existing) target time. In this way, the time may be thought of as being responsible of re-adjusting itself (in a non-linear way), once the fluctuations in the brane worlds occur, so as to restore the conformal invariance of the underlying world sheet theory, which has been disturbed by

the brane fluctuations.

One of the most important results of [9] is the appearance of a time-dependent central charge deficit in the underlying conformal theory, acting as a vacuum energy density in the respective target-space lagrangian. This also is crucial for a ‘graceful exit’ from the inflationary de Sitter phase, and the absence of eternal acceleration. In fact the asymptotic (in time) theory is that of a flat (Minkowski) target-space σ -model with a linear dilaton [6] in the string frame. In the Einstein frame, i.e. in a redefined metric background in which the Einstein curvature term in the target-space effective action has the canonical normalization, the universe is linearly expanding, which is the limiting case in which the horizon (1.5) diverges logarithmically; hence such a theory can admit properly defined asymptotic states and S -matrix amplitudes. The linear dilaton background has been shown [6] to be a consistent background for string theory, despite being time dependent, in the sense of satisfying factorizability (for certain discrete values of the asymptotic central charge though), modular invariance and unitarity.

Another important aspect of our solution is the fact that the extra bulk dimensions are compactified in such a way that one is significantly larger than the others, thereby leading to effective five-dimensional brane world scenarios. The reason for this is appropriately chosen five-form flux background. This feature is, we believe, one of the most important ones of the type-0 stringy cosmologies, which are known to be characterized by the existence of non-trivial flux form fields coming from the Ramond sector of the brane worlds [11, 9].

The reader might object to our use of type-0 backgrounds due to the existence of tachyonic backgrounds. Although at tree level it has been demonstrated that the above-described flux forms can stabilize such backgrounds, by shifting away the tachyonic mass poles, however, recently this feature has been questioned at string loop level. Nevertheless, in the context of our cosmological model, such quantum instabilities are expected probably as a result of the *non-equilibrium* nature of our relaxing background, in which the time

dependent dilaton field plays the rôle of the quintessence field. Indeed, as demonstrated in [9] the asymptotic in time value of the tachyon background is zero, and hence such a field disappears eventually from the spectrum, which is consistent with the asymptotic equilibrium nature of the ground state.

The structure of the present article, is as follows: in section 2 we review the main features of the cosmological model of [9]. In section 3 we discuss the phenomenology of the model as implied by the current astrophysical observations. In particular, we demonstrate that the present-era values of the deceleration parameter of the type-0 non-critical string Universe, the “vacuum energy density”, and the Hubble parameter all match the experimental data. A crucial rôle for this, in particular for matching the order-one value of the deceleration parameter, is played by the relative magnitude of the flux of the five form field of the type 0 string to the size of the five extra dimensions transverse to the direction of the flux. Conclusions are presented in section 4.

2. Cosmology with type-0 non-critical Strings: A brief review

In this section we review the main features of the Cosmological String model of [9]. We commence our discussion by recalling that the effective ten-dimensional target space action of the type-0 String, to $\mathcal{O}(\alpha')$ in the Regge slope α' , assumes the form [11]:

$$\begin{aligned}
 S = \int d^{10}x \sqrt{-G} & \left[e^{-2\Phi} \left(R + 4(\partial_M \Phi)^2 - \frac{1}{4}(\partial_M T)^2 - \frac{1}{4}m^2 T^2 \right. \right. \\
 & \left. \left. - \frac{1}{12}H_{MNP}^2 \right) - \frac{1}{4} \left(1 + T + \frac{T^2}{2} \right) |\mathcal{F}_{MNP\Sigma T}|^2 \right]
 \end{aligned} \tag{2.1}$$

where capital Greek letters denote ten-dimensional indices, Φ is the dilaton, H_{MNP} denotes the field strength of the antisymmetric tensor field, which we shall ignore in the present work, and T is a tachyon field of mass $m^2 < 0$. In our analysis we have ignored higher than quadratic order terms in the tachyon potential. The quantity $\mathcal{F}_{MNP\Sigma T}$ denotes the appropriate five-form of type-0 string theory, with non trivial flux, which couples to the

tachyon field in the Ramond-Ramond (RR) sector via the function $f(T) = 1 + T + \frac{1}{2}T^2$.

From (2.1) one sees easily the important rôle of the five-form $\mathcal{F}$ in stabilizing the ground state. Due to its special coupling with the quadratic T^2 term in Ramond-Ramond (RR) sector of the theory, it yields an effective mass term for the tachyon which is positive, despite the originally negative m^2 contribution [11]. As mentioned previously, such a stability has recently been questioned in the context of string loop corrections, but as we have mentioned previously this is rather a desirable feature of the approach, in view of the claimed cosmological instabilities.

As argued in [9] *fluctuations of the brane worlds* involved in the construction of type-0 string theory result in *supercriticality* of the underlying σ -model, with inevitable consequence the addition of the following term to the action (2.1)

$$- \int d^{10}x \sqrt{-G} e^{-2\Phi} Q(t)^2 \quad (2.2)$$

where Φ is the dilaton field, and $Q(t)$ is the central-charge deficit of the non-equilibrium non conformal σ -model theory. The time here is identified with the (world-sheet zero mode of) the Liouville field, and the t dependence of the central charge deficit is in accordance with the concept of a Zamolodchikov C-function [12], a “running central charge” of a non-conformal theory, interpolating between two conformal (fixed point) theories.

The sign of $Q(t)^2$ is positive if one assumes supercriticality of the string [6, 5, 10, 9], which is the case of the model of [9]. It is important to remark that in general, $Q^2(t)$ depends on the σ -model backgrounds fields, being the analogue of Zamolodchikov’s C -function [12]. As explained in [9], the explicit time dependence of $Q(t)$ reflects the existence of relevant operators in the problem, other than the background fields considered in (2.1) which are treated collectively in the present context. Such operators have been argued to represent *initial* quantum fluctuations of the brane world. A plausible scenario, for instance, would be that the initial disturbance that takes the system out of equilibrium is due to an impulse on the D3 brane worlds coming from either a scattering off it of a macroscopic number of closed

string bulk states or another brane in scenarios where the bulk space is uncompactified (e.g. ekpyrotic universes *etc.* [13]). For times long after the event, memory of the details of this process is kept in the temporal evolution of $Q^2(t)$, which is determined self-consistently by means of the Liouville equations, as we shall see below.

We note in passing that such time-, and background field, -dependent deficits also appear [14] if one views a standard critical string on a $d+1$ -dimensional cosmological string background as a *non critical* σ -model propagating on a spatially-dependent d -dimensional background. In such a case the time dependence of the spatial coordinates $x^i(t)$, $i = 1, \dots, d$, is attributed to the fact that the latter represent trajectories in the $d+1$ -dimensional space time. In this respect, the starting point is a d -dimensional non-conformal σ -model with only *spatially dependent* backgrounds. The extra time variable then is viewed as a time-like Liouville field which restores conformal invariance of the d -dimensional stringy σ -model. In this way, the model has a time-dependent (‘running’) central charge deficit, being given essentially by the d -dimensional target-space effective action. In our approach [7, 10, 9], however, as we have stressed repeatedly, the dimensionality of the target space time is not increased by the Liouville-dressing procedure. Instead, the time (=Liouville) field itself readjusts its configuration in a non-linear way in order to restore the conformal invariance broken by the fluctuations of the brane worlds in the type-0 string theory [7, 9].

The ten-dimensional metric configuration we considered in [9] was:

$$G_{MN} = \begin{pmatrix} g_{\mu\nu}^{(4)} & 0 & 0 \\ 0 & e^{2\sigma_1} & 0 \\ 0 & 0 & e^{2\sigma_2} I_{5 \times 5} \end{pmatrix} \quad (2.3)$$

where lower-case Greek indices are four-dimensional space time indices, and $I_{5 \times 5}$ denotes the 5×5 unit matrix. We have chosen two different scales for internal space. The field σ_1 sets the scale of the fifth dimension, while σ_2 parametrizes a flat five dimensional space. In the context of cosmological models, we are dealing with here, the fields $g_{\mu\nu}^{(4)}$, σ_i , $i = 1, 2$

are assumed to depend on the time t only.

As we demonstrated in [9], a consistent background choice for the flux form field will be that in which the flux is parallel to the fifth dimension σ_2 . This implies actually that the internal space is crystallized (stabilized) in such a way that this dimension is much larger than the remaining four σ_1 .

Upon considering the fields to be time dependent only, i.e. considering spherically-symmetric homogeneous backgrounds, restricting ourselves to the compactification (2.3), and assuming a Robertson-Walker form of the four-dimensional metric, with scale factor $a(t)$, the generalized conformal invariance conditions and the Curci-Pafutti σ -model renormalizability constraint [15] imply a set of differential equations, which we solved numerically in [9].

The generic form of these equations reads [5, 7, 9]:

$$\ddot{g}^i + Q(t)\dot{g}^i = -\tilde{\beta}^i \quad (2.4)$$

where $\tilde{\beta}^i$ are the Weyl anomaly coefficient of the stringy σ -model on the background $\{g^i\}$. In the model of [9] the set of $\{g^i\}$ contains graviton, dilaton, tachyon, flux and moduli fields $\sigma_{1,2}$ whose vacuum expectation values control the size of the extra dimensions. The equations, then, have the following explicit form:

$$\begin{aligned} -3\frac{\ddot{a}}{a} + \ddot{\sigma}_1 + 5\ddot{\sigma}_2 - 2\ddot{\Phi} + \dot{\sigma}_1^2 + 5\dot{\sigma}_2^2 + \frac{1}{4}\dot{T}^2 + e^{-2\sigma_1+2\Phi}f_5^2f(T) &= 0 , \\ \ddot{a}a + a\dot{a}\left(2Q + \dot{\sigma}_1 + 5\dot{\sigma}_2 - 2\dot{\Phi}\right) + e^{-2\sigma_1+2\Phi}f_5^2f(T)a^2 &= 0 , \\ \ddot{\sigma}_1 + 5\dot{\sigma}_1^2 + 3\frac{\dot{a}}{a}\dot{\sigma}_1 + 2Q\dot{\sigma}_1 + 5\dot{\sigma}_1\dot{\sigma}_2 - 2\dot{\sigma}_1\dot{\Phi} + e^{-2\sigma_1+2\Phi}f_5^2f(T) &= 0 , \\ 3\ddot{\sigma}_2 + 9\dot{\sigma}_2^2 + 3\frac{\dot{a}}{a}\dot{\sigma}_2 + 2Q\dot{\sigma}_2 + \dot{\sigma}_1\dot{\sigma}_2 - 2\dot{\sigma}_2\dot{\Phi} - e^{-2\sigma_1+2\Phi}f_5^2f(T) &= 0 , \\ 2\ddot{T} + 3\frac{\dot{a}}{a}\dot{T} + Q\dot{T} + \dot{\sigma}_1\dot{T} + 5\dot{\sigma}_2\dot{T} - 2\dot{T}\dot{\Phi} + \\ m^2T - 4e^{-2\sigma_1+2\Phi}f_5^2f'(T) &= 0 , \end{aligned}$$

$$\begin{aligned}
& \ddot{\Phi} + Q\dot{\Phi} + 6\frac{\dot{a}}{a} + 6\frac{\dot{a}^2}{a^2} + \\
& 2 \left[-\ddot{\sigma}_1 - \frac{3\dot{a}}{a}\dot{\sigma}_1 - 5\ddot{\sigma}_2 - 15\frac{\dot{a}}{a}\dot{\sigma}_2 - \dot{\sigma}_1^2 - 15\dot{\sigma}_2^2 - 5\dot{\sigma}_1\dot{\sigma}_2 - \right. \\
& \left. 2\dot{\Phi}^2 + 2\ddot{\Phi} + 6\frac{\dot{a}}{a}\dot{\Phi} + 2\dot{\sigma}_1\dot{\Phi} + 10\dot{\sigma}_2\dot{\Phi} \right] - \frac{1}{4}\dot{T}^2 + \frac{1}{4}m^2T^2 + Q^2 = 0 , \\
& C_5 = e^{-\sigma_1+5\sigma_2}f(T)f_5 , \\
& \Phi^{(3)} + Q\ddot{\Phi} + \dot{Q}\dot{\Phi} + 12\frac{\dot{a}}{a^3} \left(a\ddot{a} + \dot{a}^2 + Qa\dot{a} \right) - \dot{T}(\ddot{T} + Q\dot{T}) + \\
& 4\dot{\sigma}_1(\ddot{\sigma}_1 + 2\dot{\sigma}_1^2 + Q\dot{\sigma}_1) + 20\dot{\sigma}_2(\ddot{\sigma}_2 + 2\dot{\sigma}_2^2 + Q\dot{\sigma}_2) = 0
\end{aligned} \tag{2.5}$$

where $f'(T)$ denotes functional differentiation with respect to the field T , the overdot denotes time derivative, with respect to the σ -model frame, and $\Phi^{(3)}$ denotes triple time derivative.

As argued in [9] such equations correspond to solutions of equations of motion derived from a ten-dimensional effective action. This is an important and non-trivial consequence of the gradient flow property of the σ -model $\tilde{\beta}^i$ functions, according to which:

$$\tilde{\beta}^i = \mathcal{G}^{ij} \frac{\delta \mathcal{F}[g]}{\delta g^j} \tag{2.6}$$

where the flow functional $\mathcal{F}[g]$ is essentially the target-space effective action, depending on the background configuration under consideration.

An equivalent set of equations (in fact at most linear combinations) come out from the corresponding four-dimensional action after dimensional reduction. Of course this reduction leads to the string or σ -model frame, in which there are dilaton exponential factors in front of the Einstein term in the action. We may turn to the Einstein frame, in which such factors are absent, and the Einstein term is canonically normalized, through the transformation

$$g_E = e^{-2\Phi+\sigma_1+5\sigma_2}g \tag{2.7}$$

In this frame the line element is

$$ds_E^2 = -e^{-2\Phi+\sigma_1+5\sigma_2}dt^2 + a^2(t)e^{-2\Phi+\sigma_1+5\sigma_2}(dr^2 + r^2d\Omega^2) \tag{2.8}$$

Therefore to discuss cosmological evolution we have to pass to the cosmological time defined by

$$dt_E = e^{-\Phi + \frac{\sigma_1 + 5\sigma_2}{2}} dt \quad (2.9)$$

Then the line element, for a spatially flat universe, which we assume here motivated by the CMB data [3], becomes:

$$ds_E^2 = -dt_E^2 + a_E^2(t_E)(dr^2 + r^2 d\Omega^2) \quad (2.10)$$

with

$$a_E(t_E) = e^{-\Phi + \frac{\sigma_1 + 5\sigma_2}{2}} a(t(t_E)) . \quad (2.11)$$

Recalling the notation [9]:

$$a = e^{b(t)t} \quad (2.12)$$

in the σ -model frame, one has the following useful relations between the Einstein and σ -model frames, to be used here:

$$\begin{aligned} \frac{da_E}{dt_E} &= e^b \left(-\dot{\Phi} + \frac{\dot{\sigma}_1 + 5\dot{\sigma}_2}{2} + \dot{b} \right) , \\ \frac{d^2 a_E}{dt_E^2} &= e^b e^{\Phi - \frac{\sigma_1 + 5\sigma_2}{2}} \left[\dot{b} \left(-\dot{\Phi} + \frac{\dot{\sigma}_1 + 5\dot{\sigma}_2}{2} + \dot{b} \right) - \ddot{\Phi} + \frac{\ddot{\sigma}_1 + 5\ddot{\sigma}_2}{2} + \ddot{b} \right] . \end{aligned} \quad (2.13)$$

The Hubble parameter reads:

$$H(t_E) \equiv \frac{\frac{da_E}{dt_E}}{a_E} = e^{\Phi - \frac{\sigma_1 + 5\sigma_2}{2}} \left(-\dot{\Phi} + \frac{\dot{\sigma}_1 + 5\dot{\sigma}_2}{2} + \dot{b} \right) \quad (2.14)$$

while the deceleration parameter of the Universe (1.2) acquires the form:

$$q = - \frac{\dot{b} \left(-\dot{\Phi} + \frac{\dot{\sigma}_1 + 5\dot{\sigma}_2}{2} + \dot{b} \right) - \ddot{\Phi} + \frac{\ddot{\sigma}_1 + 5\ddot{\sigma}_2}{2} + \ddot{b}}{\left(-\dot{\Phi} + \frac{\dot{\sigma}_1 + 5\dot{\sigma}_2}{2} + \dot{b} \right)^2} \quad (2.15)$$

Finally the Einstein-frame “vacuum” energy is related to the central charge deficit [9]

$$\Lambda_E = e^{2\Phi - \sigma_1 - 5\sigma_2} Q^2(t) \quad (2.16)$$

As discussed in [9], due to the non-equilibrium nature of the non-critical string Universe, which has not yet relaxed to its ground state, Λ_E should be considered rather as an effective

potential, in much the same way as the potential of a (non-equilibrium) quintessence field, whose rôle is played here the dilaton Φ [8, 9].

We should also remark that we have adopted the usual Kaluza-Klein reduction without considering the extra dimensions compactified. Nevertheless the equations we are interested in and the results we will discuss in the following section are not modified if we had considered a compact six-dimensional space instead. In that case $e^{2\sigma_1}$ and $e^{2\sigma_2}$ should correspond to radii of the compact space. Note also that in the string frame we have the exponential $e^{-2\Phi+\sigma_1+5\sigma_2}$ instead of $e^{-2\Phi}$, since we allow time dependence of the volume of the compact space.

The numerical solution we have found is supported by analytical considerations for the asymptotic field modes (late cosmological-frame times $t \rightarrow \infty$). We followed an iterative method of solving the system of these equations. The starting point of the iteration procedure is the solution of the linear system with the correct asymptotic behaviour. Then we insert the linear solution into the system of equations and keep only the linear part to get the general solution. For details and results we refer the reader in [9].

The most important physical results of our analysis may be summarized as follows: our solution demonstrates that the scale factor of the Universe, after the initial singularity, enters a short inflationary phase and then, in a smooth way, goes into a flat Minkowski spacetime with a linear dilaton for asymptotically long σ -model times $t \rightarrow \infty$ in the σ -model frame. Equivalently, in the Einstein frame, this asymptotic string theory corresponds to a string propagating into a linearly expanding, non-accelerating Universe, with a (negative valued) dilaton that varies logarithmically with the Robertson-Walker (Einstein frame) time. Note that this is a consistent σ -model background [6] in the sense of satisfying modular invariance and factorization of S -matrix elements. This is actually one of the most important points of our work in [9], namely that there is a smooth exit from the de Sitter phase in such a way that one can *appropriately define* asymptotic on-shell states,




Figure 1: The evolution of the central charge deficit Q^2 in the Einstein frame. Immediately after inflation, the deficit passes through a phase where it first vanishes, and then oscillates before relaxing to an equilibrium constant value asymptotically.

and hence an S -matrix. The fields σ_1 and σ_2 which parametrize the internal space have an interesting behaviour. The field σ_1 , which sets the scale of the fifth dimension, during inflation contracts until it reaches a constant value. After inflation, it maintains this value, until the universe evolves to the above mentioned asymptotically flat spacetime in the σ -model frame. The field σ_2 which parametrizes the conformally flat five-dimensional space freezes to a constant value which is much smaller than that of the fifth dimension. Thus we see that, in our model, a cosmological evolution may lead to different scales for the extra dimensions. It is important to notice, that this difference in scales of the extra dimensions is due to the fact that in our theory the gravity is very weak asymptotically. A phenomenologically important feature of our model is that the vacuum energy, determined by the central-charge deficit, relaxes to zero asymptotically in a way which is reminiscent of quintessence models, with the rôle of the quintessence field played by the dilaton. During the *relaxation process* the string theory remains consistent as a conformal σ -model, since the target time, which here plays the rôle of the Liouville field, restores the broken conformal




Figure 2: The evolution of the decelerating parameter q of the type-0 string Universe in the Einstein frame.

invariance [7, 10, 9].

The Cosmology of the model in the late time phase, where the extra dimensions have been stabilized to their equilibrium values and the tachyonic mode has relaxed to a constant value (assumed zero by normalization) [9], may be summarized diagrammatically in figures 1 and 2. In these figures we give the evolution (in the Einstein frame) of the central charge deficit and the deceleration parameter with time. The figures refer to the numerical solution of [9].

From figure 1 we observe that, immediately after inflation, the central charge deficit Q^2 passes through a *metastable point* where it vanishes. This has to do with the fact that at this point the square root Q changes sign ¹, as can also be confirmed analytically from the corresponding expression for the linearized solution of [9]. After this point the central

¹This change of sign of Q is irrelevant for Liouville σ -model physics, although important for target space physics as we shall discuss later on. Indeed, in the original Liouville σ -model Lagrangian one encounters only Q^2 factors [5]. To obtain a canonical normalization for the Liouville σ -model kinetic term one needs only to perform the redefinition [5] $\phi \rightarrow |Q(t)|\phi$ to the Liouville mode, which is thus unaffected by a sign change of $Q(t)$, with t the world-sheet zero mode of the Liouville field ϕ .

charge deficit oscillates (as a function of time) before it relaxes to its constant asymptotic value, which should be one of the values for which the conformal theory of [6] is valid. The oscillatory nature is consistent with the time-like signature of the Liouville mode as explained in [9]. This behaviour of the central charge deficit is important in determining the evolution of the energy densities of the type-0 string Universe, as we shall discuss later on.

We also observe from fig. 2 that, after its graceful exit from the inflationary phase, the type-0 non-critical string Universe passes first through a decelerating phase, which is then succeeded by an accelerating one, before the Universe relaxes asymptotically to its steady state (equilibrium) value.

It is the purpose of this article to discuss the precise phenomenology of this Universe, and compare it with current observational evidence [1]. To this end we shall compute the value of the deceleration parameter at the “present era”. The concept of ‘present era’ will be defined appropriately in section 3.

Before doing so, we would like to make an important comment concerning the initial singularity which characterizes our solution. The singularity is a general feature of the equations of the form (2.5). The singularity may be a true one, in case one thinks of the initial brane fluctuation which caused the non-criticality as a *catastrophic event* (e.g. a situation in the ekpyrotic universe scenario [13]), or it may be removable, in case the initial fluctuation is a quantum fluctuation in the context of the M-theory. In the latter case, removing the singularity is probably a matter of a full quantum description of the theory, which at present is not available. We note at this stage, however, the possibility of deriving smooth cosmological solutions in string theory, without initial singularities, by including in the action higher curvature terms (e.g. quadratic of Gauss-Bonnet type [16]), which are part of the quantum corrections, in the sense of being generated by including string-loop corrections in the effective action. For our purposes here, such issues are not

directly relevant.

3. Present Era Phenomenology of the non-critical type-0 String Universe: Compatibility with supernovae Ia observations

Having obtained numerical evidence (c.f. fig. 2) from the full numerical solution on the existence of an accelerating phase of the type-0 string Universe, before its final relaxation into the critical string regime, we now proceed to study *analytically* this phase. In particular, in this section, we shall be interested in studying the *precise* behaviour, as well as estimating the *order of magnitude*, of the *deceleration parameter* q (2.15) for Einstein frame times long after the initial fluctuation of the $D3$ brane worlds. This would allow comparison with the current observations.

In this late time regime the various fields can be approximated well by their *linearized solution*, described in detail in [9]. Recalling that for the Einstein-frame time one has

$$t_E = t_{E,0} + \int_{t_0}^t d\tau \left(e^{-\Phi(\tau) + \frac{\sigma_1(\tau) + 5\sigma_2(\tau)}{2}} \right) \quad (3.1)$$

we mention that, for the numerical solution of [9], the various field modes can be approximated sufficiently well by the linear solution for $t_0 \geq 25$ in string units of time.

We will also assume that at present times, long after the inflationary period, which we are interested in here, the extra dimensions, as well as the tachyon field, are already frozen to their constant asymptotic values $\sigma_{1,0} \equiv s_{01}$, $\sigma_{2,0} \equiv s_{02}$. The validity of this assumption will be discussed later on. For the tachyon field, this freezing-out behaviour implies that $f(T) = 1 + T + T^2/2 \rightarrow 1$, while the freezing out of the extra dimensions implies that the “volume” $V_6 \equiv e^{s_{01} + 5s_{02}}$ of the extra dimensions can be absorbed, as usual after compactification, in the four-dimensional gravitational coupling constant. This will be assumed in what follows, which means that, from now on, the string and the Einstein times in (3.1) will be related only through the dilaton field, without explicit involvement of V_6 .

The freezing out of the extra dimensions and the tachyon to their constant (equilibrium) values implies that any non-trivial dependence on the fields σ_1, σ_2 and T disappears from the equations satisfied by the other fields.

Keeping, therefore, only the modes a, Q, Φ, f_5 in the remaining equations (2.5), we observe that they acquire the form:

$$\begin{aligned}
& \frac{C_5^2 e^{2\sigma_{01}}}{2V_6^2} e^{2\Phi} - 3\frac{\ddot{a}}{a} - 2\ddot{\Phi} = 0, \\
& \frac{C_5^2 e^{2\sigma_{01}}}{2V_6^2} e^{2\Phi} + (2Q - 2\dot{\Phi})\frac{\dot{a}}{a} + \frac{\ddot{a}}{a} = 0, \\
& \Phi^{(3)} + Q\ddot{\Phi} + \dot{Q}\dot{\Phi} = 0 \quad (\text{Curci - Paffuti relation}), \\
& f_5 = \frac{C_5 e^{2s_{01}}}{V_6} \quad (\text{five - form equation}), \\
& -Q^2 - Q\dot{\Phi} + 4\dot{\Phi}^2 - 6\frac{\ddot{a}}{a} - 12\frac{\dot{a}}{a}\dot{\Phi} - 5\ddot{\Phi} = 0 \quad (\text{dilaton equation}). \tag{3.2}
\end{aligned}$$

Note that the five-form field is also frozen to a constant value. In the Curci-Paffuti relation we have dropped out the combination $12\frac{\dot{a}}{a^3}(a\ddot{a} + \dot{a}^2 + Qa\dot{a})$, which is very small compared to the other terms, for the time period we are interested in.

The Curci-Paffuti relation in (3.2) can be integrated for the field Q and the solution respecting the asymptotic behaviour $Q \rightarrow q_0$ is:

$$Q = -\frac{F_1 q_0}{\dot{\Phi}} - \frac{\ddot{\Phi}}{\dot{\Phi}} \tag{3.3}$$

where F_1 is a positive constant and $\dot{\Phi} \rightarrow -F_1$ (asymptotically linear dilaton). The system of the remaining three non-trivial equations then reads:

$$\begin{aligned}
& \alpha_1 + 0 \times \frac{\dot{a}}{a} - 3\frac{\ddot{a}}{a} = 0, \\
& \beta_1 + (2Q - 2\dot{\Phi})\frac{\dot{a}}{a} + \frac{\ddot{a}}{a} = 0, \\
& \gamma_1 - 12\dot{\Phi}\frac{\dot{a}}{a} - 6\frac{\ddot{a}}{a} = 0, \\
& \text{where } \alpha_1 \equiv \frac{C_5^2}{2V_6^2 e^{-2s_{01}}} e^{2\Phi} - 2\ddot{\Phi}, \quad \beta_1 \equiv \frac{C_5^2}{2V_6^2 e^{-2s_{01}}} e^{2\Phi},
\end{aligned}$$

$$\gamma_1 \equiv -Q^2 - Q\dot{\Phi} + 4\dot{\Phi}^2 - 5\ddot{\Phi} \quad (3.4)$$

Compatibility requires the vanishing of the determinant

$$0 = \det \begin{pmatrix} \alpha_1 & 0 & -3 \\ \beta_1 & 2Q - 2\dot{\Phi} & 1 \\ \gamma_1 & -12\dot{\Phi} & -6 \end{pmatrix} \quad (3.5)$$

which yields an equation for Φ , whose solution, by compatibility, must yield the linear solution $\Phi \simeq f_0 - F_1 t$ for large times t . Indeed, this is what happens, as we shall demonstrate below.

We first observe that to linear order in the fields one obtains:

$$Q = q_0 + \frac{q_0}{F_1}(F_1 + \dot{\Phi}). \quad (3.6)$$

From this we, therefore, see that $Q \rightarrow q_0$ if and only if $\dot{\Phi} \rightarrow -F_1$, which, as we shall see below, is true.

Indeed, from the resolvent (3.5), keeping only terms linear in the fields, we obtain for $\Phi(t)$:

$$\ddot{\Phi}(t) + \alpha^2 A^2 e^{2\Phi(t)} + \beta^2 F_1 (F_1 + \dot{\Phi}) = 0 \quad (3.7)$$

where $\alpha^2 = \frac{11+\sqrt{17}}{2(3+\sqrt{17})}$, $\beta^2 = \frac{17+5\sqrt{17}}{3+\sqrt{17}}$ are numerical constants of order $\mathcal{O}(1)$, and $A^2 \equiv \frac{C_5^2 e^{2s_{01}}}{2V_6^2}$.

For times long after the initial fluctuations, such as the present times, where the linear approximation is valid, the term $\beta^2 F_1 (F_1 + \dot{\Phi})$ is hierarchically small, and may be neglected in the dilaton equation (3.7), yielding $\alpha^2 A^2 e^{2\Phi} + \ddot{\Phi} \simeq 0$. This is solved for:

$$\Phi(t) = -\ln \left[\frac{\alpha A}{F_1} \cosh(F_1 t) \right], \quad (3.8)$$

with F_1 a positive constant. For large times $F_1 t \gg 1$ (in string units) one therefore recovers the linear solution for the dilaton, thereby demonstrating the self-consistency of the approach: $\Phi \sim f_0 - F_1 t$, $F_1 = |f_1| > 0$ ².

²From (3.8) we thus observe that the asymptotic weakness of gravity in this Universe [9] is due to the

Defining the Einstein frame time t_E as

$$t_E = \int^t e^{-\Phi(z)} dz \quad (3.9)$$

we get

$$t_E = \frac{\alpha A}{F_1^2} \sinh(F_1 t). \quad (3.10)$$

The string frame time t , can be expressed in terms of t_E as:

$$t = \frac{1}{F_1} \ln \left[\sqrt{1 + \frac{F_1^4}{\alpha^2 A^2} t_E^2} + \frac{F_1^2}{\alpha A} t_E \right]. \quad (3.11)$$

In terms of the Einstein-frame time (3.9) one obtains a logarithmic time-dependence [6] for the dilaton

$$\Phi_E = \text{const} - \ln t_E, \quad (3.12)$$

For this behaviour of Φ , the central charge deficit (3.6) tends to a constant value q_0 . This value must be, for consistency of the underlying string theory, *one of the discrete values* obtained in [6], for which the factorization property (unitarity) of the string scattering amplitudes occurs. Notice that this asymptotic string theory, with a constant (time independent) central-charge deficit, $Q^2 \propto c - 25$ (or $c - 9$ for superstring) is considered an *equilibrium* situation, where an S -matrix can be defined for specific (discrete) values of the central charge c . The standard critical (super)string corresponds to central charge $c = 25$ ($=9$ for superstrings) [5, 6].

We would like at this point to go back for a moment and justify the validity of the assumption that the extra dimensions are frozen, and hence any non-trivial dependence on the fields $\sigma_{1,2}$ can be ignored. The evolution of the extra dimensions is determined by the third and forth of the equations (2.5)

$$A^2 e^{2\Phi} - 2\dot{\sigma}_1 \left(\dot{\Phi} + \frac{\ddot{\Phi}}{\dot{\Phi}} + \frac{q_0 F_1}{\dot{\Phi}} \right) + \dot{\sigma}_1 \dot{\sigma}_2 + 5\dot{\sigma}_1^2 + \ddot{\sigma}_1 = 0$$

smallness of the internal space V_6 as compared with the flux C_5 of the five form field, $f_0 \sim \ln V_6$ (c.f. previous expression for A). We shall come back to this important point later on.

$$-A^2 e^{2\Phi} - 2\dot{\sigma}_2 \left(\dot{\Phi} + \frac{\ddot{\Phi}}{\dot{\Phi}} + \frac{q_0 F_1}{\dot{\Phi}} \right) + \dot{\sigma}_1 \dot{\sigma}_2 + 9\dot{\sigma}_2^2 + 3\ddot{\sigma}_2 = 0 \quad (3.13)$$

In the range where the linear approximation is valid this system takes the form

$$\begin{aligned} \ddot{\sigma}_1 + 2(q_0 + F_1)\dot{\sigma}_1 + \frac{F_1^2}{\alpha^2 \cosh^2(F_1 t)} &= 0 , \\ 3\ddot{\sigma}_2 + 2(q_0 + F_1)\dot{\sigma}_2 - \frac{F_1^2}{\alpha^2 \cosh^2(F_1 t)} &= 0 . \end{aligned} \quad (3.14)$$

The above system has indeed regular solutions of the type assumed above, namely solutions that freeze out to constant values relatively quickly after inflation. To see this, let us concentrate for definiteness to the first of the equations (3.14). The second equation can be analysed in a formally identical way.

Integrating the first of equations (3.14) we obtain

$$\dot{\sigma}_1 + 2(q_0 + F_1)\sigma_1 + \frac{F_1}{\alpha^2} \tanh(F_1 t) = c_1, \quad c_1 = \text{integration constant} .$$

Weighting this equation by $e^{2(q_0+F_1)t}$ and integrating once more we obtain:

$$\sigma_1 + \frac{1}{\alpha^2} \text{ncosh}(F_1 t) - \frac{1}{\alpha^2} e^{-2(q_0+F_1)t} \int dt' e^{2(q_0+F_1)t'} \text{ncosh}(F_1 t') = \frac{c_1}{2(q_0 + F_1)} + e^{-2(q_0+F_1)t} c_2 , \quad (3.15)$$

where $c_{1,2}$ are integration constants. We shall be interested in the regime of long times $F_1 t \gg 1$ (in string units) after inflation. In this regime the t -integration can be done by approximating $\text{ncosh}(F_1 t) \simeq F_1 t$. After some straightforward algebra we then obtain:

$$\sigma_1 \rightarrow s_{01} = \frac{c_1/F_1 - 1}{2(1 + q_0/F_1)\alpha^2} + \mathcal{O}\left(e^{-2(q_0+F_1)t} c_2\right) \quad (3.16)$$

where $q_0, F_1 > 0$. The reader should bear in mind that the solution of [9] requires

$$\left(\frac{q_0}{F_1} \right) = \frac{1}{2}(1 + \sqrt{17}) \simeq 2.56 . \quad (3.17)$$

This relation stems from the dilaton equation, which guarantees that the Liouville-dressed theory preserves conformal invariance, as discussed in detail in [9].

In a similar way one finds that the field σ_2 asymptotes quickly to

$$\sigma_2 \rightarrow s_{02} = \frac{c'_1/F_1 + 1}{2(1 + q_0/F_1)\alpha^2} + \mathcal{O}\left(e^{-\frac{2}{3}(q_0+F_1)t}c'_2\right) \quad (3.18)$$

where $c'_{1,2}$ are appropriate integration constants.

The constants in the equilibrium values s_{0i} , $i = 1, 2$ can be chosen in such a way [9] that the size of the σ_1 dimension is larger than the rest. We also remind the reader at this point that in this regime of large string-frame times $F_1 t \gg 1$ the dilaton can be approximated by its linear form $\Phi \sim -F_1 t$. This completes our digression on the self-consistency of the approximation of ignoring the effects of the σ_i fields for large times $F_1 t \gg 1$.

We now come to discuss the behaviour of the scale factor $a(t)$. From the second of (3.2) one obtains the evolution equation for the scale factor,

$$\frac{\ddot{a}}{a} + (2q_0 + F_1)\frac{\dot{a}}{a} + \frac{F_1^2}{\alpha^2} \frac{1}{\cosh^2(F_1 t)} = 0, \quad (3.19)$$

This can be easily solved by means of the Gauss hypergeometric functions ${}_2F_1$; in the Einstein frame one has:

$$\begin{aligned} a_E(t_E) = & \frac{F_1}{\gamma} (\sqrt{1 + \gamma^2 t_E^2}) \left[C_1 \left(\frac{\sqrt{1 + \gamma^2 t_E^2}}{\gamma t_E + \sqrt{1 + \gamma^2 t_E^2}} \right)^{\frac{F_1 + q_0}{F_1}} \times \right. \\ & {}_2F_1 \left(\frac{1}{4} \left(1 - \frac{2(F_1 + q_0)}{F_1} - \frac{\sqrt{4 + \alpha^2}}{\alpha} \right), \right. \\ & \left. \frac{-2\alpha(F_1 + q_0) + F_1(\alpha + \sqrt{4 + \alpha^2})}{4F_1\alpha}, 1 - \frac{F_1 + q_0}{F_1}, \frac{1}{1 + \gamma^2 t_E^2} \right) + \\ & C_2 \left[\sqrt{1 + \gamma^2 t_E^2} + \gamma t_E \right]^{-\frac{F_1 + q_0}{F_1}} {}_2F_1 \left(\frac{1}{4} \left(1 + \frac{2(F_1 + q_0)}{F_1} - \frac{\sqrt{4 + \alpha^2}}{\alpha} \right), \right. \\ & \left. \frac{2\alpha(F_1 + q_0) + F_1(\alpha + \sqrt{4 + \alpha^2})}{4F_1\alpha}, 1 + \frac{F_1 + q_0}{F_1}, \frac{1}{1 + \gamma^2 t_E^2} \right) \left. \right] \end{aligned} \quad (3.20)$$

where $C_{1,2}$ are integration constants, and

$$\gamma \equiv \frac{F_1^2}{\alpha A}, \quad A \equiv \frac{|C_5|e^{s_{01}}}{V_6} = |C_5|e^{-5s_{02}} \quad (3.21)$$

Notice the independence of A on the large compact dimension s_{01} . This will play an important physical rôle as we shall discuss later on.

For large t_E , e.g. present cosmological time values, one has

$$a_E(t_E) \simeq \frac{F_1}{\gamma} \sqrt{1 + \gamma^2 t_E^2} \quad (3.22)$$

For very large (future) times $a(t_E)$ scales linearly with the Einstein-frame cosmological time t_E [9], and hence the cosmic horizon (1.5) disappears, thereby allowing the proper definition of asymptotic states and thus a scattering matrix. Asymptotically therefore in time, the Universe relaxes to its ground-state equilibrium situation, and the non-criticality of the string, caused by the initial fluctuation, disappears, making room for a critical (equilibrium) string Universe.

The Hubble parameter (2.14) reads for large t_E

$$H(t_E) \simeq \frac{\gamma^2 t_E}{1 + \gamma^2 t_E^2} \quad (3.23)$$

while the deceleration parameter (1.2) in the same regime of t_E becomes:

$$q(t_E) \simeq -\frac{1}{\gamma^2 t_E^2} \quad (3.24)$$

Finally, the “vacuum energy” reads:

$$\Lambda(t_E) \simeq \frac{q_0^2 \gamma^2}{F_1^2 (1 + \gamma^2 t_E^2)} \quad (3.25)$$

From (3.23), (3.25) we observe that one can match the present-era values quite straightforwardly, as expected by naive dimensional analysis. On the other hand, the dimensionless deceleration parameter q (3.24), although negative, appears to be extremely suppressed as compared to the order one value inferred from the best fit of the supernova data (1.1),(1.3). We should remark, however, that, due to the current uncertainties in the data [17], this is not necessarily in contradiction with the current observations for a non-zero cosmological constant. Nevertheless, if one takes the best fit Universe (1.1) literally, then it seems that

in order to match all the data one should obtain an order one deceleration parameter of the type 0 string Universe (c.f. discussion following (1.3)).

In our case this is possible, since as we observe from (3.20), in the Einstein frame the scale factor of our type-0 non-critical string Universe is only a function of the combination γt_E . If, therefore, one defines the *present era* by the time regime

$$\gamma \sim t_E^{-1} \quad (3.26)$$

in the Einstein frame, then from (3.20) it becomes clear that an order one negative value of q is obtained.

The important point, however, is that this is compatible with large enough times t_E (in string units) for

$$|C_5|e^{-5s_{02}} \gg 1 . \quad (3.27)$$

as becomes clear from the definition of γ (3.21). This condition can be guaranteed *either* for small radii of the five of the extra dimensions *or* for a large value of the flux $|C_5|$ of the five-form of the type-0 string. Notice that the relatively large extra dimension, in the direction of the flux, s_{01} , decouples from this condition, thus allowing for the possibility of effective five-dimensional models with large uncompactified fifth dimension.

Notice that in the regime (3.26) of Einstein-frame times the Hubble parameter and the cosmological constant will continue to be compatible with the current observations, and in fact to depend on $\gamma \sim t_E^{-1}$ as in their large γt_E regime given above (3.23),(3.25). This was expected from simple dimensional analysis and can be confirmed by a detailed analysis of the respective formulae obtained from (3.20).

We next look at the equation of state of our type-0 string Universe. As discussed in [9], our situation is a quintessence like case, with the dilaton playing the rôle of the quintessence field [8, 9]. Hence the equation of state for our type-0 string Universe reads [2]:

$$w_\Phi = \frac{p_\Phi}{\rho_\Phi} = \frac{\frac{1}{2}(\dot{\Phi})^2 - V(\Phi)}{\frac{1}{2}(\dot{\Phi})^2 + V(\Phi)} \quad (3.28)$$

where p_Φ is the pressure and ρ_Φ is the energy density, and $V(\Phi)$ is the effective potential for the dilaton, which in our case is provided by the central-charge deficit term. Here the dot denotes Einstein-frame differentiation.

In the Einstein frame the potential $V(\Phi)$ is given by Λ_E in (3.25). In the limit $Q \rightarrow q_0$, which has been argued to characterize the present era to a good approximation, the present era $V(\Phi)$ is then of order $(q_0^2/2F_1^2)t_E^{-2}$, where we recall that q_0/F_1 is given by (3.17).

In the Einstein frame the exact normalization of the dilaton field is $\Phi_E = \text{const} - \ln t_E$. Combining this result with (3.17), we then obtain for the present era (3.26):

$$\frac{1}{2}\dot{\Phi}^2 \sim \frac{1}{2t_E^2}, \quad V(\Phi) \sim \frac{6.56}{2} \frac{1}{t_E^2} \quad (3.29)$$

This implies an equation of state (3.28):

$$w_\Phi(t_E \gg 1) \simeq -0.74 \quad (3.30)$$

for (large) times t_E in string units corresponding to the present era (3.26).

We should compare this result with the case of an “honest” cosmological *constant* situation, as the one fitting the data in [1], which yields $w = -1$, or with the standard scenario of a *slow-varying quintessence field* φ , $\dot{\varphi}^2 \ll V(\varphi)$, which again yields $w_\Phi \simeq -1$ by means of (3.28). The reader should recall that, in our case, the result (3.30) occurs because the relative magnitude (3.29) of the dilaton (quintessence) potential versus that of its kinetic energy is fixed by conformal invariance (3.17). It would be interesting to see whether the inclusion of ordinary matter (attached on the 3-branes of the type-0 string) changes these results significantly. This is left for future work.

If the present era of the Universe is therefore defined by (3.26), then one obtains *compatibility* of the above results with the *current astrophysical observations* [1, 3]:

$$H(t_E) \sim \frac{1}{t_E}, \quad \Lambda(t_E) \sim \frac{\mathcal{O}(1-10)}{t_E^2} > 0, \quad q(t_E) \sim -|\mathcal{O}(1)| < 0 \quad (3.31)$$

It is amusing to see here that in order to get an order-one deceleration parameter (3.31), as the present phenomenology suggests [1], it appears necessary to have the ratio of the flux C_5 of the five-form of the type 0 string over the volume of the (five) transverse dimensions $e^{5\sigma_{02}}$ very large. If one insists on keeping the flux C_5 of order one, something, however, which is *not necessary*, then this result implies *transplanckian* (smaller than Planck length) sizes of at least five of the extra dimensions, so that the volume $V_6 \ll 1$ in string units.

Such small volumes in turn require bulk string mass scales much larger than the Planck mass scale on the four-dimensional brane world. To see this recall that the gravitational part of the bulk ten-dimensional target-space effective action (2.1) reads after compactification in the Einstein frame:

$$S \sim \frac{1}{g_{10}^2} M_s^8 V_6 \int d^4x \sqrt{g^{(4)}} R_E^{(4)} + \dots \quad (3.32)$$

where $g_{10} \sim e^{\langle \Phi \rangle}$ is the ten-dimensional string coupling, which for our purposes here is assumed weak $g_{10} < 1$. The overall coefficient in front of the four-dimensional Einstein term defines the square of the four dimensional Planck scale M_P :

$$M_P^2 = \frac{1}{g_{10}^2} M_s^8 V_6 \quad (3.33)$$

Therefore, for small $V_6 \ll 1$ in string units, i.e. $M_s^6 V_6 \ll 1$, one obtains that the string mass scale should be much larger than the four-dimensional Planck mass scale:

$$M_P \ll M_s . \quad (3.34)$$

Notice that, in the modern viewpoint that the string bulk scale is a free parameter in string/M-theory, such relations are allowed. This is a curious feature of our construction, and certainly requires further study; in particular, one should investigate possible connections between our approach and that of [18], where transplanckian modes have been argued to play an important rôle for inflation.

Notice that the above results have been obtained without the inclusion of ordinary matter. In the type-0 string/brane scenario, ordinary matter may be assumed attached to

the brane, and hence purely four dimensional. The latter is going to resist the deceleration of the universe, according to standard arguments, but it is not expected to change the order of magnitude of the above quantities. In this sense one may obtain the observed ‘coincidence situation’ of the present era, where the matter and ‘dark energy’ contributions are roughly of the same order of magnitude [2]. In our scenario it is the time dependence of both ‘dark energy’ and ‘matter contribution’, in conjunction with the value the time t_E has at present, roughly $t_E \sim 10^{60} M_P^{-1}$, that is held responsible for this coincidence situation.




Figure 3: The evolution of the energy densities of matter, radiation and of the quintessence field (dilaton) vs. the scale factor of the Universe in the Einstein frame. At early stages the energy density of the quintessence field decreases significantly, as compared with the rest, and the coincidence situation is lost. This is due to the behaviour of the central charge deficit of the model, shown in figure 1, which dives in to zero for a short period immediately after inflation.

Indeed, from fig. 1 we observe that at relatively early times (after inflation) there is a period where the central charge deficit Q^2 , and hence the potential energy $V(\Phi)$ of the quintessence ‘tracking’ dilaton field Φ , vanishes. The dilaton equation, which guarantees conformal invariance of the Liouville dressed theory [9], implies near this point that

$(d\Phi/dt_\sigma)^2 \sim d^2\Phi/dt_\sigma^2$, where $t_\sigma = -\ln t_E$ is the σ -model time. This means that the dilaton field kinetic energy density, and hence its total energy density ρ_Λ , scales logarithmically with the Einstein time in this (short) time region $\rho_\Lambda \propto 1/(\ln t_E + \text{const})^2$. Shortly after this point, as the time elapses the central charge increases significantly in such a way that $d^2\Phi/dt^2 < 0$, and eventually the dilaton reaches its linear equilibrium configuration in string frame. This behaviour is dictated by conformal invariance of the underlying σ -model. Taking into account (3.8),(3.9), this implies that, as one goes backwards in time, starting from the present era, the energy density of Φ , $\rho_\Phi = \frac{1}{2}(\dot{\Phi})^2 + V(\Phi)$, becomes significantly smaller than the energy density of matter (or radiation), which increase as the time goes backward, scaling with the scale factor like a_E^{-3} (or a_E^{-4}). Thus in our model the tracking (coincidence) of the matter energy density by that of the quintessence dilaton field is a feature only of the present era, which is a welcome feature phenomenologically. The situation is summarized in figure 3, where we plot (in a qualitative manner) the energy densities of radiation, matter and of the quintessence field (dilaton), ρ_Λ , vs. the scale factor $a_E(t_E)$ of the Robertson-Walker non-critical string Universe (in the Einstein frame). The plot, which is not to scale, is based on the (qualitative) behaviour of the Q^2 , shown in figure 1, and the above discussion.

As the time t_E elapses, the matter contribution will become subdominant, as scaling like a_E^{-3} . For very large times t_E in the far future, as we have seen above, the dominant contributions will be the ones due to the non-constant in time ‘dark energy component’ $\Lambda(t_E) \sim a_E^{-2}$, which asymptotes to zero, as the system reaches its equilibrium value. This makes a quantitative difference in scaling as compared with the standard Robertson-Walker scenario with a constant vacuum energy (c.f. (1.4)).

4. Conclusions

In this note we have discussed the cosmological evolution of the present era deceleration parameter of a non-critical type-0 string Universe, and we have argued that one can get compatibility with current astrophysical observations. We have been able to fit the data with a quintessence-like non-critical string Universe, which has a present-era negative deceleration parameter q of order one, in agreement with the supernova Ia observations. The dark energy of our string Universe, however, is not constant, but relaxes to zero asymptotically, in a way compatible with the current value of the dark energy, explaining the observed ‘coincidence’ in the order of magnitudes between matter and dark energy components as a matter of ‘chance’ (this is like an anthropic principle situation: we have been ‘lucky’ to witness this event, being in the right ‘place’ at the ‘right time’).

We do not claim here that our crude string model is a physical model for the Universe, but we find it interesting that at least to a first approximation the phenomenology of the model seems to match the data. An important rôle for obtaining an order-one value for the deceleration parameter is played by the relative value of the flux of the five-form field of the type-0 string to the volume of the small extra dimensions. Interestingly enough the value of the deceleration parameter is independent of the large bulk dimension, along the direction of which lies the flux of the five form of the type-0 string. This leaves room for accommodating effective five-dimensional scenaria, with large uncompactified fifth dimension.

We therefore consider these considerations very interesting, and certainly worthy of further investigations. It is our belief that these characteristics extend beyond the specific model of type-0 string theory studied here. In fact we think that such a behaviour may characterize a large class of non-critical (non-equilibrium) string Universes, relaxing to their critical situation asymptotically in time. It will certainly be interesting to study properly the phenomenology of such models, by taking into account the detailed form of string matter, attached to the brane worlds, including fermionic excitations, and see whether

interesting predictions can be made, in the cosmological sense, that could differentiate among the various models. We hope to come back to such a study in the near future.

Acknowledgements

G.A.D and B.C.G. would like to acknowledge partial financial support from the Athens University special account for research. N.E.M. wishes to thank Subir Sarkar for discussions. E.P. wishes to thank the Physics Department of King’s College London for the warm hospitality during the last stages of this work.

References

- [1] A. G. Riess *et al.*, *Astroph. J.* **117**, 707 (1999); S. Perlmutter *et al.*, *Astroph. J.* **517**, 565 (1999).
- [2] For a comprehensive and up to date review see: S. M. Carroll, *Living Rev. Rel.* **4**, 1 (2001) [arXiv:astro-ph/0004075].
- [3] J.R. Bond, A.H. Jaffe and L. Knox, *Phys. Rev.* **D57**, 2117 (1998).
- [4] A.P. Billyard, A.A. Coley and J.E. Lidsey, *J. Math. Phys.* **41** 6277 (2000); S. Hellerman, N. Kaloper and L. Susskind, [hep-th/0101180]; W. Fischler, A. Kashani-Poor, R. McNees, S. Paban, [hep-th/0104181]; E. Witten, hep-th/0106109 ; P. O. Mazur and E. Mottola, *Phys. Rev. D* **64**, 104022 (2001) [arXiv:hep-th/0106151], and references therein.
- [5] F. David, *Mod. Phys. Lett.* **A3**, 1651 (1988); J. Distler and H. Kawai, *Nucl. Phys.* **B321**, 509 (1989); see also N.E. Mavromatos and J.L. Miramontes, *Mod. Phys. Lett.* **A4**, 1847 (1989); E. D’Hoker and P.S. Kurzepa, *Mod. Phys. Lett.* **A5**, 1411 (1990); E. D’Hoker, *Mod. Phys. Lett.* **A6**, 745 (1991); for earlier references see: E.D’Hoker

- and R. Jackiw, Phys. Rev. **D26**, 3517 (1982); A. Bilal and J.L. Gervais, Nucl. Phys. **B284**, 397 (1987); J.L. Gervais and A. Neveu, Nucl. Phys. **B224**, 329 (1983).
- [6] I. Antoniadis, C. Bachas, John Ellis and D.V. Nanopoulos, Phys. Lett. **B211** 383 (1988); Nucl. Phys. **B328** 117 (1989).
- [7] J. Ellis, N. E. Mavromatos and D. V. Nanopoulos, Phys. Lett. **B293**, 37, (1992); Mod. Phys. Lett. **A10** (1995) 1685; for reviews see: Erice Subnuclear Series (World Sci., Singapore) **31** 1, (1993); [hep-th/9304133]; J. Chaos, Solitons and Fractals **10**, 345 (eds. C. Castro and M.S. El Naschie, Elsevier Science, Pergamon 1999) [hep-th/9805120].
- [8] J. R. Ellis, N. E. Mavromatos and D. V. Nanopoulos, hep-th/0105206.
- [9] G. A. Diamandis, B. C. Georgalas, N. E. Mavromatos, E. Papantonopoulos and I. Pappa, arXiv:hep-th/0107124, Int. J. Mod. Phys. **A** in press.
- [10] G. A. Diamandis, B. C. Georgalas, N. E. Mavromatos and E. Papantonopoulos, Phys. Lett. **B461**, 57 (1999) [hep-th/9903045].
- [11] I. Klebanov and A.A. Tseytlin, Nucl. Phys. **B546**, 155 (1999); Nucl. Phys. **B547**, 143 (1999).
- [12] A.B. Zamolodchikov, JETP Lett. **43** (1986) 730; Sov. J. Nucl. Phys. **46** (1987) 1090.
- [13] J. Khoury, B. A. Ovrut, P. J. Steinhardt and N. Turok, Phys. Rev. D **64**, 123522 (2001) [arXiv:hep-th/0103239].
- [14] C. Schmidhuber and A.A. Tseytlin, Nucl. Phys. **B426**, 187, (1994).
- [15] G. Curci and G. Paffuti, Nucl. Phys. **B286**, 399, (1987).
- [16] See, for instance: I. Antoniadis, J. Rizos and K. Tamvakis, Nucl. Phys. B **415**, 497 (1994) [hep-th/9305025].

- [17] see, for instance: Subir Sarkar, JHEP Conference Proceedings, *Budapest 2001, High energy physics hep2001/299*, [arXiv:hep-ph/0201140].
- [18] J. Martin and R. H. Brandenberger, Phys. Rev. D **63**, 123501 (2001) [arXiv:hep-th/0005209].